

CLINIC-SPECIFIC DEMOGRAPHIC, CLINICAL, AND LABORATORY DATA

The remainder of this report provides clinic-specific figures for each of the 26 currently participating clinics. Individual figures for each clinic show demographic and clinical characteristics of the men with gonorrhea enrolled in GISP, as well as antimicrobial susceptibilities for the *N. gonorrhoeae* isolates. The number of isolates submitted by each clinic is 300 when the full sample of 25 isolates per month is obtained. However, the number of isolates submitted is lower for many clinics located in areas with low gonorrhea rates. Each page of figures is labeled with the city of the participating clinic and the actual number of isolates on which the clinic's 1999 data are based.

Definitions of terms and abbreviations used in the clinic-specific figures are given below.

Figure D: contact=has sexual partner with gonorrhea

Figure F: ceftriaxone 250=ceftriaxone 250 mg
 ceftriaxone 125=ceftriaxone 125 mg
 other cephalo=other cephalosporins

Figure G: azi/ery=azithromycin/erythromycin
 doxy/tet=doxycycline/tetracycline

Figure H: PPNG=penicillinase-producing *N. gonorrhoeae*
 TRNG=plasmid-mediated tetracycline resistant *N. gonorrhoeae*
 PPNG-TRNG=plasmid-mediated penicillin and tetracycline resistant *N. gonorrhoeae*
 PenR=chromosomally mediated penicillin resistant *N. gonorrhoeae*
 TetR=chromosomally mediated tetracycline resistant *N. gonorrhoeae*
 CMRNG=chromosomally mediated penicillin and tetracycline resistant *N. gonorrhoeae*

Figure K: intermediate res=intermediate resistance